


Sensitivity of Coastal Environments and Wildlife to Spilled Oil NORTHERN CALIFORNIA



Office of
OIL SPILL PREVENTION and RESPONSE
California Department Of Fish and Game
Sacramento, California

Prepared for:



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration
Seattle, Washington


Prepared by:



RESEARCH PLANNING, INC.
Columbia, SC 29202


SEPTEMBER 1994

ENVIRONMENTAL SENSITIVITY INDEX MAP




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration



PUBLISHED: SEPTEMBER 1994


DRAKES BAY, CALIF. (1976)

NC-1

ENVIRONMENTAL SENSITIVITY INDEX MAP

122°52'30"

122°45'00"



122°52'30"

122°45'00"




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration




PUBLISHED: SEPTEMBER 1994

POINT REYES NE, CALIF. (1976)
INVERNESS, CALIF. (1976)


NC-2

ENVIRONMENTAL SENSITIVITY INDEX MAP




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for




HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration



PUBLISHED: SEPTEMBER 1994


TOMALES, CALIF. (1976) **NC-3**

ENVIRONMENTAL SENSITIVITY INDEX MAP




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for




HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration



PUBLISHED: SEPTEMBER 1994


VALLEY FORD, CALIF. (1971) **NC-4**

ENVIRONMENTAL SENSITIVITY INDEX MAP




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration




PUBLISHED: SEPTEMBER 1994

BODEGA HEAD, CALIF. (1972)

NC-5

ENVIRONMENTAL SENSITIVITY INDEX MAP




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration



PUBLISHED: SEPTEMBER 1994


DUNCANS MILLS, CALIF.(1979)

NC-6

ENVIRONMENTAL SENSITIVITY INDEX MAP

123°15'00"

123°07'30"



123°15'00"

123°07'30"




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration




PUBLISHED: SEPTEMBER 1994

FORT ROSS, CALIF. (1978)
ARCHED ROCK, CALIF. (1977)


NC-7

ENVIRONMENTAL SENSITIVITY INDEX MAP




Office of
OIL SPILL PREVENTION and RESPONSE
California Department of Fish and Game

Prepared for



HAZARDOUS MATERIALS RESPONSE
and ASSESSMENT DIVISION
National Oceanic and Atmospheric Administration



PUBLISHED: SEPTEMBER 1994

PLANTATION, CALIF. (1977)





















NC-8

NORTHERN CALIFORNIA

SHORELINE HABITAT RANKINGS
































- 1A EXPOSED ROCKY CLIFFS
- 1B EXPOSED SEAWALL
- 1C EXPOSED ROCKY CLIFFS WITH BOULDER TALUS BASE
- 2 WAVE CUT ROCKY PLATFORMS
- 3 FINE- TO MEDIUM-GRAINED SAND BEACHES
- 4 COARSE-GRAINED SAND TO GRANULE BEACHES
- 5 MIXED SAND AND GRAVEL BEACHES
- 6A GRAVEL BEACHES
- 6B RIPRAP
- 7 EXPOSED TIDAL FLATS
- 8A SHELTERED ROCKY SHORES
- 8B SHELTERED MAN-MADE STRUCTURES
- 9 SHELTERED TIDAL FLATS
- 10A SALT MARSHES

HUMAN-USE FEATURES

-  ACCESS POINT
-  AIRPORT
-  AQUACULTURE
-  ARCHAEOLOGICAL SITE
-  BOAT RAMP
-  COAST GUARD
-  COMMERCIAL FISHING
-  HISTORICAL SITE
-  HOIST
-  MARINA
-  MARINE SANCTUARY
-  NATIONAL PARK
-  STATE PARK
-  RECREATIONAL FISHING AND BOATING
-  RECREATIONAL BEACH
-  SUBSISTENCE AREA
-  VILLAGE
-  WATER INTAKE
-  WILDLIFE REFUGE
-  ID NUMBER

----- PARK BOUNDARY

SENSITIVE BIOLOGICAL RESOURCES

- | | | |
|--|--|---|
| <p> BIRDS</p> <ul style="list-style-type: none">  ALCIDS & PELAGIC BIRDS  DIVING BIRDS  GULLS & TERNS  RAPTORS  SHOREBIRDS  WADING BIRDS  WATERFOWL  NESTING SITES | <p> MAMMALS</p> <ul style="list-style-type: none">  DOLPHINS  MUSTELIDS & RODENTS  SEA OTTERS  SEALS & SEA LIONS  WHALES | <p> SHELLFISH</p> <ul style="list-style-type: none">  BIVALVES  CRABS  ECHINODERMS  GASTROPODS & ABALONE  LOBSTERS  SHRIMP  SQUID |
| <p> FISH</p> <ul style="list-style-type: none">  FISH | <p> PLANTS</p> <ul style="list-style-type: none">  KELP & SEAGRASSES  TERRESTRIAL PLANTS | <p> MULTI-GROUP</p> <ul style="list-style-type: none">  THREATENED/ENDANGERED  ID NUMBER |

Guidelines for Interpreting ESI Maps

To help users interpret the ESI maps and tabular data, we offer the following guidelines for use in addition to the map legend:

- **Shoreline Habitats.** The “shoreline,” representing the boundary between land and water, is color-coded with the ESI classification. Most shoreline habitats are shown as a line, with no areal dimension. Where there is more than one shoreline type (e.g., a beach in front of a seawall), the colors for each habitat are shown, with the color for the landward habitat on the land side of the shoreline and the color for the seaward habitat on the water side. In areas where the intertidal zone is wide (e.g., wide tidal flats, wave-cut rocky platforms), the habitat from high to low water is filled with the ESI classification color. When data are available, the entire extent of wetlands are filled with colored patterns. The seaward edge of the wetland is color-coded with the ESI classification; the landward extent of the wetland is indicated by a dashed, colored line.
- **Biological Resources.** The distribution of biological resources is shown using many different conventions. The major convention is an icon associated with a point, line, or polygon that shows the species’ areal distribution. The icon’s reference number corresponds to a data table with details on species and life history. Biological resource data are organized into six major groups, each with a reference color: birds (green), mammals (brown), fish (blue), shellfish (orange), reptiles (red), and rare/endangered plants and special habitats (purple). These colors are used to fill hatched polygons and the icons. Each major group has subgroups with unique icons to visually indicate the type of organism or feature present. The icon or group of icons is usually located inside the polygon it represents; however, sometimes a line is connected between the icon and the polygon or point to make it easier to relate the two. Note that icons are used to indicate the types of resources present, but the actual data are the points and polygons. A red box around an icon indicates the presence of a species on the state or Federal list of threatened or endangered species.

The number listed below each icon refers to the first column of a data table for each map. The data tables, organized by group (birds, fish, etc.), include the following information: species name, status as threatened or endangered on state and Federal lists, concentration (specifically for each point or polygon), presence by month, and special life-history time periods. When a polygon contains multiple groups, the one number under the group of icons is listed under each group heading in the data tables. Where possible, the same number is used on multiple maps. For example, all bald eagle nests with the same seasonality could have the same number throughout the atlas, or the same assemblage of fish would have the same number wherever it occurred.

A data table has a separate listing for every unique combination of species, concentration, seasonality, life-history stage, and source. By looking at the monthly seasonality data in the table for each map, the species present at the time of concern can be easily identified. An ‘X’ or number is placed under each month in which any life stage of the species is present in the area represented by the point or polygon. Numbers are used typically for fish and shellfish where data on relative abundance are available. The final columns in the data tables include the months when reproductive activities occur or early life stages are present. Users should pay close attention to the data tables because they contain much of the information needed to identify the most sensitive resources at different times of the year.

Points, lines, and polygons on a map represent the distribution of the resources. Green points show bird nesting sites, including bald eagle nests and dense colonial nesters (e.g., heron rookeries and seabird nesting colonies). Animals and habitats are also represented as: 1) hatched polygons in the color for the animal group (e.g., green for birds); 2) black hatched polygons which contain multiple groups of resources (birds and fish in the same tidal channels); 3) solid lines (usually used for fish in small streams); or 4) in “common in ...” boxes. When showing the biological resource polygons would make the maps too difficult to read (usually when multiple polygons cover a large area), the polygons are not plotted and the presence of the resource is indicated by placing the icon in a box labeled “common in ...” The box contains an appropriate geographic reference. Different boxes can be used on the same map when, for example: “common in Winyah Bay” or “common in tidal creeks.” The data for these resources are still fully present in the database but are not shown to make the maps more readable.

- **Human-use Resources.** Most of the human-use resources are point features indicated by a black-and-white icon. Managed lands, such as refuges and sanctuaries, have their boundaries shown as a dot-dash line with an icon and name placed inside. Where the feature is a known point location (e.g., a drinking water intake, boat ramp, marina), the exact location is shown as a small black dot and a line is drawn from it to the icon. Activities such as commercial and recreational fishing and areas such as recreational beaches are also indicated by an icon placed in the general area without any lines to points or polygons since the boundaries are not readily defined.

Some features, like historic and archaeological sites, are location-sensitive: the agency managing the resource believes the exact location should not be shown in order to protect the site. In these cases, the icon is placed in the general area of the resource, but the exact location is not shown.

ESIMAP 1 (CONTINUED)

BIOLOGICAL RESOURCES:

BIRD:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
	Red-breasted merganser				X	X	X	X					X	X	X	X	-	-	-	-
	Red-necked grebe				X	X	X								X	X	-	-	-	-
	Sanderling																-	-	-	-
	Short-billed dowitcher																-	-	-	-
	Sora rail				X	X	X	X					X	X	X	X	-	-	-	-
	Surf scoter				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Virginia rail				X	X	X	X	X	X	X	X	X	X	X	X	APR-AUG	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Western gull				X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	White-winged scoter				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Willet				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
209	Brown pelican	F	E		X	X	X	X	X	X	X	X	X	X	X	X	FEB-NOV	FEB-JUL	MAR-AUG	JUN-NOV
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Peregrine falcon	F	E		X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Sooty shearwater						X	X	X	X	X	X	X	X	X	X	-	-	-	-
210	Common murre				X	X	X	X	X	X	X	X	X	X	X	X	MAY-SEP	MAY-JUL	JUN-AUG	JUL-SEP
211	Black scoter (common)				X	X	X	X						X	X	X	-	-	-	-
	Common loon				X	X	X	X					X	X	X	X	-	-	-	-
	Eared grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Harlequin duck				X	X	X	X					X	X	X	X	-	-	-	-
	Horned grebe				X	X	X	X	X	X			X	X	X	X	-	-	-	-
	Pied-billed grebe				X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-APR	APR-JUL	JUN-SEP
	Red-breasted merganser				X	X	X	X					X	X	X	X	-	-	-	-
	Red-necked grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Surf scoter				X	X	X	X					X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
	White-winged scoter				X	X	X	X					X	X	X	X	-	-	-	-
212	Black scoter (common)				X	X	X	X						X	X	X	-	-	-	-
	Common loon				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Eared grebe																-	-	-	-
	Harlequin duck				X	X	X	X					X	X	X	X	-	-	-	-
	Horned grebe				X	X	X	X	X				X	X	X	X	-	-	-	-
	Marbled godwit				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Pied-billed grebe				X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-APR	APR-JUL	JUN-SEP
	Red-breasted merganser				X	X	X	X					X	X	X	X	-	-	-	-
	Red-necked grebe				X	X	X							X	X	-	-	-	-	-
	Sanderling																-	-	-	-
	Surf scoter				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Western gull				X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	White-winged scoter				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Willet				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
213	Canvasback				X	X	X							X	X	X	-	-	-	-
	Northern phalarope				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Northern shoveler																-	-	-	-
	Red phalarope				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
558	Black oystercatcher			6	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Brandt's cormorant			1522	X	X	X	X	X	X	X	X	X	X	X	X	FEB-AUG	FEB-JUN	MAR-JUL	MAY-AUG
	Common murre			15155	X	X	X	X	X	X	X	X	X	X	X	X	MAY-SEP	MAY-JUL	JUN-AUG	JUL-SEP
	Pelagic cormorant			266	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			616	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Rhinoceros auklet			6	X	X	X	X	X	X	X	X	X	X	X	X	MAY-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Tufted puffin			4	X		X	X	X	X	X	X	X	X	X	X	APR-NOV	APR-AUG	MAY-SEP	JUL-NOV
	Western gull			178	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
730	Black scoter (common)				X	X	X	X						X	X	X	-	-	-	-
	Brown pelican	F	E		X	X	X	X	X	X	X	X	X	X	X	X	FEB-NOV	FEB-JUL	MAR-AUG	JUN-NOV
	Common loon				X	X	X	X					X	X	X	X	-	-	-	-
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Eared grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Harlequin duck				X	X	X	X					X	X	X	X	-	-	-	-
	Horned grebe				X	X	X	X	X				X	X	X	X	-	-	-	-
	Peregrine falcon	F	E		X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Pied-billed grebe				X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-APR	APR-JUL	JUN-SEP
	Red-breasted merganser				X	X	X	X					X	X	X	X	-	-	-	-
	Red-necked grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Sooty shearwater						X	X	X	X	X	X	X	X	X	X	-	-	-	-
	Surf scoter				X	X	X	X					X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
	White-winged scoter				X	X	X	X					X	X	X	X	-	-	-	-

FISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Outmigration
197	Pacific herring			HIGH	X	X	X							X	X	X	OCT-MAR	-
199	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-
200	Pacific herring				X	X	X							X	X	X	OCT-MAR	-
201	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-
207	Pacific herring				X	X	X							X	X	X	OCT-MAR	-
209	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-
730	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-

M_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
20	Harbor seal			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN
198	Dall's porpoise				X	X				X	X	X				X	-	-	-
	Gray whale			HIGH	X	X	X	X						X	X	-	-	-	-
	Harbor porpoise				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-
199	California sea lion			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAY-AUG
	Northern (Steller) sea lion	F	T	MED						X	X	X	X			-	-	-	MAY-AUG
	Northern elephant seal			LOW	X	X	X				X	X			X	-	-	-	DEC-MAR
201	California sea lion			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAY-AUG
	Harbor seal			LOW	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN
	Northern (Steller) sea lion	F	T	MED						X	X	X	X			-	-	-	MAY-AUG

ESIMAP 1 (CONTINUED)

BIOLOGICAL RESOURCES:

M_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
	Northern elephant seal			MED	X	X	X					X	X			X	-			DEC-MAR
203	Harbor seal			MED	X	X	X	X	X	X	X	X	X	X	X	X	-			MAR-JUN
205	Minke whale			MED						X	X	X	X	X	X					-
208	Northern elephant seal									X	X	X								-
209	California sea lion			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-			MAY-AUG
	Harbor seal			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-			MAR-JUN
	Northern (Steller) sea lion	F	T	HIGH					X	X	X	X								MAY-AUG
	Northern elephant seal			HIGH	X	X	X			X	X					X	-			DEC-MAR
730	California sea lion			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-			MAY-AUG
	Northern (Steller) sea lion	F	T	MED					X	X	X	X								MAY-AUG
	Northern elephant seal			LOW	X	X	X			X	X					X	-			DEC-MAR

PLANT:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
60	Salt marsh bird's-beak	F	E													
197	Eelgrass			HIGH												
199	Bull kelp															
200	Eelgrass			HIGH												
201	Bull kelp															
202	Beach layia	F	E													
	Clover lupine	F	E													
207	Eelgrass			HIGH												
209	Bull kelp															
730	Bull kelp															

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Spawning
197	Common Pacific littleneck clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP
	Gaper clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-
198	Dungeness crab			HIGH	X	X	X	X	X							X	-	-
199	Abalone																-	-
	Red sea urchin				X	X	X	X	X	X	X	X	X	X	X	X	-	-
200	Bay ghost shrimp				X	X	X	X	X	X	X	X	X	X	X	X	-	JUN-JUL
	Gaper clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-
201	Abalone																-	-
	Red sea urchin				X	X	X	X	X	X	X	X	X	X	X	X	-	-
207	Bay ghost shrimp				X	X	X	X	X	X	X	X	X	X	X	X	-	JUN-JUL
	Gaper clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-
209	Abalone																-	-
	Red sea urchin				X	X	X	X	X	X	X	X	X	X	X	X	-	-
730	Abalone																-	-
	Red sea urchin				X	X	X	X	X	X	X	X	X	X	X	X	-	-

HUMAN USE RESOURCES:

AQUACULTURE:

RAR#	Name	Owner	Contact	Phone
H202	M-438-01 PARCEL 2-JOHNSON OYSTER CO.	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H203	M-438-01 PARCEL 1-JOHNSON OYSTER CO.	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521

HISTORICAL_SITE:

RAR#	Name	Owner	Contact	Phone
H215	LIFE SAVING STATION			

ESIMAP 2

BIOLOGICAL RESOURCES:

BIRD:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
197	American avocet				X	X	X	X					X	X	X	X	-	-	-	-
	Canvasback			HIGH	X	X	X						X	X	X	-	-	-	-	
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Greater scaup			HIGH	X	X	X	X					X	X	-	-	-	-	-	
	Marbled godwit				X	X	X	X				X	X	X	X	-	-	-	-	
	Osprey			LOW	X	X	X	X	X	X	X	X	X	X	X	X	MAR-APR	APR-JUN	MAY-JUL	JUL-AUG
	Peregrine falcon	F	E	HIGH	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	
	Western grebe				X	X	X	X				X	X	X	X	-	-	-	-	
	Willet				X	X	X	X				X	X	X	X	-	-	-	-	
214	Canvasback				X	X	X						X	X	X	-	-	-	-	
	Common loon				X	X	X	X	X				X	X	X	-	-	-	-	
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Greater scaup				X	X	X	X					X	X	-	-	-	-	-	
	Western grebe				X	X	X	X				X	X	X	X	-	-	-	-	
216	American avocet				X	X	X	X				X	X	X	X	-	-	-	-	
	California black rail	S	T	HIGH	X	X	X	X	X	X	X	X	X	X	X	X	MAR-JUN	MAR-JUN	-	-
	Great blue heron				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Great egret				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Marbled godwit				X	X	X	X				X	X	X	X	-	-	-	-	
	Peregrine falcon	F	E	HIGH	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	
	Willet				X	X	X	X				X	X	X	X	-	-	-	-	
217	Osprey			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	MAR-APR	APR-JUN	MAY-JUL	JUL-AUG

FISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Outmigration
197	Pacific herring			HIGH	X	X	X						X	X	X	OCT-MAR	-	
215	Coho salmon (silver)			HIGH	X	X	X	X	X				X	X	NOV-JAN	FEB-APR		
	Rainbow trout (steelhead)			HIGH	X	X	X	X	X				X	X	NOV-JAN	FEB-APR		

PLANT:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
60	Salt marsh bird's-beak	F	E													
197	Eelgrass			HIGH												
217	Salt marsh bird's-beak	F	E	HIGH												
218	Salt marsh bird's-beak	F	E	HIGH												

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Spawning
197	Common Pacific littleneck clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP
	Gaper clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-

T_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
215	River otter			HIGH	X	X	X	X	X	X	X	X	X	X	X	X
216	River otter			HIGH	X	X	X	X	X	X	X	X	X	X	X	X

HUMAN USE RESOURCES:

AQUACULTURE:

RAR#	Name	Owner	Contact	Phone
H192	M-430-06 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H193	M-430-09 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H194	M-430-18 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H195	M-430-13 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H196	M-430-12 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H222	M-430-05	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H223	M-430-12	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521

MARINA:

RAR#	Name	Owner	Contact	Phone
H197	MARCONI COVE MARINA	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521

ESIMAP 3 (CONTINUED)

BIOLOGICAL RESOURCES:

BIRD:

Table listing bird species (RAR#, Species), their status (S/F, T/E, Concen), and seasonal distribution (J-F, M-A, M-J, J-A, S-O, N-D, Nesting, Laying, Hatching, Fledging).

FISH:

Table listing fish species (RAR#, Species), their status (S/F, T/E, Concen), and seasonal distribution (J-F, M-A, M-J, J-A, S-O, N-D, Spawning, Outmigration).

M_MAMMAL:

Table listing mammal species (RAR#, Species), their status (S/F, T/E, Concen), and seasonal distribution (J-F, M-A, M-J, J-A, S-O, N-D, Mating, Calving, Pupping).

PLANT:

Table listing plant species (RAR#, Species), their status (S/F, T/E, Concen), and seasonal distribution (J-F, M-A, M-J, J-A, S-O, N-D).

SHELLFISH:

Table listing shellfish species (RAR#, Species), their status (S/F, T/E, Concen), and seasonal distribution (J-F, M-A, M-J, J-A, S-O, N-D, Mating, Spawning).

=====

ESIMAP 3 (CONTINUED)

HUMAN USE RESOURCES:

AQUACULTURE:

RAR#	Name	Owner	Contact	Phone
H183	M-430-15 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H184	M-430-16 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H185	M-430-17 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H186	M-430-02 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H187	M-430-11 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H188	M-430-12 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H189	M-430-08 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H190	M-430-14 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521
H191	M-430-07 AQUACULTURE LEASE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521

ESIMAP 4

BIOLOGICAL RESOURCES:

BIRD:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
176	American avocet				X	X	X	X					X	X	X	X	-	-	-	-
	Brown pelican	F	E		X	X	X	X	X	X	X	X	X	X	X	X	JAN-JUN	FEB-FEB	MAR-MAR	MAY-JUN
	Canvasback				X	X	X						X	X	X	-	-	-	-	
	Common loon				X	X	X	X	X				X	X	X	X	-	-	-	-
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Great blue heron				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Great egret				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Marbled godwit				X	X	X	X					X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Willet				X	X	X	X					X	X	X	X	-	-	-	-
177	American avocet				X	X	X	X					X	X	X	X	-	-	-	-
	Brandt's cormorant				X	X	X	X	X	X	X	X	X	X	X	X	APR-JUN	MAY-MAY	JUN-JUN	JUN-JUL
	Brown pelican	F	E		X	X	X	X	X	X	X	X	X	X	X	X	JAN-JUN	FEB-FEB	MAR-MAR	MAY-JUN
	Canvasback				X	X	X						X	X	X	-	-	-	-	
	Common loon				X	X	X	X					X	X	X	X	-	-	-	-
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Great blue heron				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Great egret				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Marbled godwit				X	X	X	X					X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
	Willet				X	X	X	X					X	X	X	X	-	-	-	-
522	Black oystercatcher			3	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Brandt's cormorant			18	X	X	X	X	X	X	X	X	X	X	X	X	FEB-AUG	FEB-JUN	MAR-JUL	MAY-AUG
	Double-crested cormorant			14	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Pelagic cormorant			143	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			20	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			32	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
547	Black oystercatcher			5	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Brandt's cormorant			12	X	X	X	X	X	X	X	X	X	X	X	X	FEB-AUG	FEB-JUN	MAR-JUL	MAY-AUG
	Pelagic cormorant			84	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			25	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			14	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP

FISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Outmigration
175	Pacific herring				X	X	X							X	X	X	OCT-MAR	-
176	Pacific herring			HIGH	X	X	X							X	X	X	OCT-MAR	-
	Rainbow trout (steelhead)			LOW	X	X	X	X						X	X	X	NOV-JAN	FEB-APR
	Striped bass			LOW	X	X	X	X	X	X	X	X	X	X	X	X	APR-JUN	-
	Tidewater goby			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	AUG-NOV	-
177	Striped bass			LOW	X	X	X	X	X	X	X	X	X	X	X	X	APR-JUN	-
	Tidewater goby				X	X	X	X	X	X	X	X	X	X	X	X	AUG-NOV	-

M_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
97	Harbor seal			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN

PLANT:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
176	Eelgrass			HIGH	-	-	-	-	-	-	-	-	-	-	-	-

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Spawning
175	Dungeness crab				X	X	X	X	X	X				X	-	-	-	-
176	Dungeness crab				X	X	X	X	X	X				X	-	-	-	-
	Gaper clam			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR
	Washington butter clam			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	-
177	Dungeness crab				X	X	X	X	X	X				X	-	-	-	-

HUMAN USE RESOURCES:

ESIMAP 5 (CONTINUED)

BIOLOGICAL RESOURCES:

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
161	Common Pacific littleneck clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP		
162	Common Pacific littleneck clam			LOW	X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP		
	Gaper clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR		
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-		
163	Gaper clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR		
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-		
164	Common Pacific littleneck clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP		
	Gaper clam			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	FEB-APR		
	Washington butter clam			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-		
166	Abalone																-	-		
167	Abalone																-	-		
170	Common Pacific littleneck clam				X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP		
171	Common Pacific littleneck clam			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	APR-SEP		
172	Dungeness crab				X	X	X	X	X	X						X	-	-		

T_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
171	River otter			HIGH	X	X	X	X	X	X	X	X	X	X	X	X

HUMAN USE RESOURCES:

AQUACULTURE:

RAR#	Name	Owner	Contact	Phone
H177	BODEGA MARINE LAB	UNIVERSITY OF CALIFORNIA	PETER CONNERS, RESERVE MANAGER	707-875-2020

MARINA:

RAR#	Name	Owner	Contact	Phone
H171	SPUD POINT MARINA	SONOMA COUNTY REGIONAL PARKS	ROBERT BLACK, SUPERVISOR	707-875-3535
H172	MASONS MARINA	MASONS MARINA	GREG ZEDRICK, MANAGER	707-875-3811
H175	PORTO BODEGA HARBOR			

WATER_INTAKE:

RAR#	Name	Owner	Contact	Phone
H173	BODEGA MARINE LAB WATER INTAKE	UNIVERSITY OF CALIFORNIA	PETER CONNERS, RESERVE MANAGER	707-875-2020
H176	BODEGA FISH FARMS - WATER INTAKE	CALIFORNIA DEPT. OF FISH AND GAME	TOM MOORE, BIOLOGIST	707-875-2521

ESIMAP 6

BIOLOGICAL RESOURCES:

BIRD:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
156	Marbled godwit				X	X	X	X					X	X	X	X	-	-	-	-
	Willet				X	X	X	X					X	X	X	X	-	-	-	-
157	Brown pelican	F	E	HIGH	X	X	X	X	X	X	X	X	X	X	X	X	JAN-JUN	FEB-FEB	MAR-MAR	MAY-JUN
	Canvasback				X	X	X						X	X	X		-	-	-	-
	Common loon				X	X	X	X	X				X	X	X		-	-	-	-
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Great blue heron				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Great egret				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Osprey			MED	X	X	X	X	X	X	X	X	X	X	X	X	MAR-APR	APR-JUN	MAY-JUL	JUL-AUG
	Red-breasted merganser				X	X	X	X					X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
541	Black oystercatcher			1	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			40	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pelagic cormorant			44	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			6	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
542	Black oystercatcher			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Brandt's cormorant			650	X	X	X	X	X	X	X	X	X	X	X	X	FEB-AUG	FEB-JUN	MAR-JUL	MAY-AUG
	Leach's storm-petrel			10	X	X	X	X	X	X	X	X	X				FEB-OCT	-	-	-
	Pelagic cormorant			44	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			34	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
543	Black oystercatcher			20	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			55	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			5	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			88	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
544	Black oystercatcher			4	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			136	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			4	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			34	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
550	Black oystercatcher			1	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Western gull			6	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP

FISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Outmigration		
155	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-		
156	Surf smelt								X	X	X	X					MAY-AUG	-		
157	Chinook salmon (king)			LOW	X	X	X	X	X								X	X	NOV-JAN	FEB-APR
	Coho salmon (silver)			HIGH	X	X	X	X									X	X	NOV-JAN	FEB-APR
	Pacific herring			MED	X	X	X										X	X	OCT-MAR	-
	Rainbow trout (steelhead)			HIGH	X	X	X	X	X								X	X	NOV-JAN	FEB-APR
	Striped bass			LOW	X	X	X	X	X	X	X	X	X	X	X	X	X	APR-JUN	-	-
	White sturgeon				X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-

M_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
157	Harbor seal			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Spawning
142	Dungeness crab			MED	X	X	X	X	X	X							X	-
155	Abalone			HIGH													-	-
	Dungeness crab			MED	X	X	X	X	X	X							X	-
	Red sea urchin			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-
157	Dungeness crab			MED	X	X	X	X	X	X							X	-

HUMAN USE RESOURCES:

ESIMAP 7

BIOLOGICAL RESOURCES:

BIRD:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
151	Marbled godwit			MED	X	X	X	X					X	X	X	X	-	-	-	-
	Willet			MED	X	X	X	X					X	X	X	X	-	-	-	-
152	Canvasback				X	X	X							X	X	X	-	-	-	-
	Common loon				X	X	X	X	X				X	X	X	X	-	-	-	-
	Double-crested cormorant				X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Great blue heron				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Great egret				X	X	X	X	X	X	X	X	X	X	X	X	MAY-JUL	MAY-JUN	JUN-JUL	JUL-AUG
	Osprey				X	X	X	X	X	X	X	X	X	X	X	X	MAR-APR	APR-JUN	MAY-JUL	JUL-AUG
	Red-breasted merganser				X	X	X	X					X	X	X	X	-	-	-	-
	Western grebe				X	X	X	X					X	X	X	X	-	-	-	-
519	Black oystercatcher			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Brandt's cormorant			51	X	X	X	X	X	X	X	X	X	X	X	X	FEB-AUG	FEB-JUN	MAR-JUL	MAY-AUG
	Double-crested cormorant			422	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	MAR-JUN	APR-JUL	MAY-AUG
	Pelagic cormorant			125	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			5	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			44	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
520	Black oystercatcher			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Brandt's cormorant			717	X	X	X	X	X	X	X	X	X	X	X	X	FEB-AUG	FEB-JUN	MAR-JUL	MAY-AUG
	Pelagic cormorant			9	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Tufted puffin			2	X		X	X	X	X	X	X	X	X	X	X	APR-NOV	APR-AUG	MAY-SEP	JUL-NOV
	Western gull			34	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
540	Black oystercatcher			7	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			227	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			20	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			42	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
541	Black oystercatcher			1	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			40	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pelagic cormorant			44	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			6	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP

FISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Outmigration	
145	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-	
150	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-	
151	Surf smelt								X	X	X	X					MAY-AUG	-	
152	Chinook salmon (king)			LOW	X	X	X	X	X							X	X	NOV-JAN	FEB-APR
	Coho salmon (silver)			HIGH	X	X	X	X	X							X	X	NOV-JAN	FEB-APR
	Rainbow trout (steelhead)			HIGH	X	X	X	X	X							X	X	NOV-JAN	FEB-APR
	White sturgeon				X	X	X	X	X	X	X	X	X	X	X	X	-	-	
153	Coho salmon (silver)			LOW	X	X	X	X	X							X	X	NOV-JAN	FEB-APR
	Rainbow trout (steelhead)			LOW	X	X	X	X	X							X	X	NOV-JAN	FEB-APR
733	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-	
734	Rockfish				X	X	X	X	X	X	X	X	X	X	X	X	-	-	

M_MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
20	Harbor seal			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN
151	Harbor seal			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN
	Northern elephant seal			LOW	X	X	X				X	X				X	-	-	DEC-MAR
152	Harbor seal			MED	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN
154	California sea lion			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAY-AUG

PLANT:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
137	Giant kelp			MED												
734	Giant kelp			MED												

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Spawning
150	Abalone			HIGH													-	-
152	Dungeness crab				X	X	X	X	X	X						X	-	-
733	Abalone			HIGH													-	-
	Red sea urchin			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-
734	Abalone			HIGH													-	-
	Red sea urchin			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-

HUMAN USE RESOURCES:

HISTORICAL_SITE:

RAR#	Name	Owner	Contact	Phone
H159	FORT ROSS STATE HISTORICAL PARK	CALIFORNIA DEPT. OF PARKS AND RECREATION	BRIAN HICKEY, CHIEF RANGER	707-865-2391

ESIMAP 8

BIOLOGICAL RESOURCES:

BIRD:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Nesting	Laying	Hatching	Fledging
480	Pelagic cormorant			121	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
536	Black oystercatcher			14	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			79	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
537	Black oystercatcher			6	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			110	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			10	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			16	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
538	Black oystercatcher			2	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			62	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Pigeon guillemot			10	X	X	X	X	X	X	X	X	X	X	X	X	MAR-AUG	APR-JUN	MAY-JUL	JUN-AUG
	Western gull			32	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
539	Black oystercatcher			1	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	MAY-JUN	JUN-JUL	JUL-SEP
	Pelagic cormorant			20	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP
	Western gull			52	X	X	X	X	X	X	X	X	X	X	X	X	MAR-SEP	APR-JUN	MAY-JUL	JUL-SEP

FISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Outmigration		
146	Rockfish			HIGH	X	X	X	X	X	X	X	X	X	X	X	X	-	-		
148	Rainbow trout (steelhead)			LOW	X	X	X	X	X								X	X	NOV-JAN	FEB-APR

M MAMMAL:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
143	Harbor seal			LOW	X	X	X	X	X	X	X	X	X	X	X	X	-	-	MAR-JUN
149	Northern (Steller) sea lion	F	T							X	X	X	X				-	-	MAY-AUG

PLANT:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D
144	Giant kelp			LOW												

SHELLFISH:

RAR#	Species	S/F	T/E	Concen	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Spawning	
146	Abalone			HIGH													-	-	
147	Dungeness crab				X	X	X	X	X	X							X	-	-

HUMAN USE RESOURCES:

HISTORICAL_SITE:

RAR#	Name	Owner	Contact	Phone
H155	FORT ROSS STATE HISTORIC PARK	CALIFORNIA DEPT. OF PARKS AND RECREATION	BRIAN HICKEY, CHIEF RANGER	707-865-2391

ENVIRONMENTAL SENSITIVITY INDEX: NORTHERN CALIFORNIA

INTRODUCTION

Environmental Sensitivity Index (ESI) maps have been developed for the shoreline of northern California to encompass the coastal areas from the Oregon border to Point Reyes. The ESI maps include information for three main components: shoreline habitats; sensitive biological resources; and human-use resources. The methods of data collection and presentation are summarized in the following sections.

SHORELINE HABITAT MAPPING

The intertidal habitats of northern California, which cover the shoreline between Point Reyes to the Oregon border, were mapped during aerial overflights and ground surveys conducted from 20-26 October 1992. The aerial surveys were conducted using the California Department of Fish and Game Partenavia, flying at elevations of 300-500 feet and slow air speed. Mapping was restricted to two hours on either side of low tides during daylight hours. An experienced coastal geologist delineated the intertidal habitats directly onto 1:24,000 scale U.S. Geological Survey topographic maps. Where appropriate, multiple habitats were delineated for each shoreline segment.

Ground teams established a total of 41 stations at sites representative of all intertidal habitats. The stations were established to ground-truth the aerial mapping. The stations also characterized the key components of each shoreline habitat in terms of the sediment distribution patterns, degree of exposure to wave and tidal current energy, dominant intertidal biota, likely oil behavior and persistence, and feasible shoreline cleanup strategies. The stations will also allow for monitoring of the seasonal trends in sediment distribution and erosional/depositional cycles, which are very important in northern California.

Prediction of the behavior and persistence of oil on intertidal habitats is based on an understanding of the dynamics of the coastal environments, not just the substrate type and grain size. The vulnerability of a particular intertidal habitat is an integration of the following factors:

- 1) Shoreline type (substrate, grain size, tidal elevation, origin)
- 2) Exposure to wave and tidal energy
- 3) Biological productivity and sensitivity
- 4) Ease of cleanup

All of these factors are used to determine the relative sensitivity of intertidal habitats. Key to the sensitivity ranking is an understanding of the relationships between: physical processes, substrate, shoreline type, product type, fate and effect, and sediment transport patterns. The intensity of energy expended upon a shoreline by wave action, tidal currents, and river currents directly affects the persistence of stranded oil. The need for shoreline cleanup activities is determined, in part, by the slowness of natural processes in removal of oil stranded on the shoreline.

These concepts have been used in the development of the Environmental Sensitivity Index (ESI), which ranks shoreline environments as to their relative sensitivity to oil spills, potential biological injury, and ease of cleanup. Generally speaking, areas exposed to high levels of physical energy, such as wave action and tidal currents, and low biological activity rank low on the scale, whereas sheltered areas with associated high biological activity have the highest ranking. The list below includes the shoreline habitats delineated for northern California, presented in order of increasing sensitivity to spilled oil.

- 1A. Exposed Rocky Cliffs
- 1B. Exposed Seawalls
- 1C. Exposed Rocky Cliffs with Boulder Talus
2. Exposed Wave-cut Platforms
3. Fine- to Medium-grained Sand Beaches
4. Coarse-grained Sand to Granule Beaches
5. Mixed Sand and Gravel Beaches
- 6A. Gravel Beaches
- 6B. Riprap
7. Exposed Tidal Flats
- 8A. Sheltered Rocky Shores
- 8B. Sheltered Man-made Structures
9. Sheltered Tidal Flats
10. Marshes

Each of the shoreline habitats are described in the following pages, in terms of their physical description, predicted oil behavior, and response considerations. Summary statistics are given for each shoreline habitat, in terms of the percent of the total shoreline length as mapped in northern California. These statistics were calculated by summing the shoreline lengths for each habitat type, double counting the segments where more than one shoreline type was mapped. Therefore, even though the length of actual shoreline mapped, which includes bays and the lower parts of rivers, was determined to 1032 kilometers, the sum of all classified shorelines was 1268 kilometers.

SENSITIVE BIOLOGICAL RESOURCES

California Department of Fish & Game (CDF&G) regional biologists in the Office of Oil Spill Prevention and Response (OSPR) compiled the biological information presented on the maps. Information collected and depicted on the maps denotes the key biological resources that are most likely at risk in the event of an oil spill. Six major categories of biological resources were considered during production of the maps: birds, fish, shellfish, mammals, plants, and reptiles. Reptiles were not included on the final copy of the maps because the only reptiles present were not likely to be impacted by a coastal oil spill or cleanup operations.

Spatial distribution of the species on the maps is represented by polygons, lines, and points, as appropriate. Associated with each of these representations is an icon depicting the types of plants or animals that are present in the polygon. Species have been divided into groups and subgroups, based on their behavior and taxonomic classification. The icons reflect this grouping scheme. The groups are color coded, and the subgroups are represented by different icons as follows:

BIRDS

-  Alcids and Pelagic Birds
-  Diving Birds
-  Gulls and Terns
-  Raptors
-  Shorebirds
-  Wading Birds
-  Waterfowl

MAMMALS

-  Dolphins
-  Mustelids and Rodents
-  Sea Otters
-  Seals and Sea Lions
-  Whales



FISH

-  Fish

SHELLFISH

-  Bivalves
-  Crabs
-  Echinoderms
-  Gastropods and Abalone
-  Lobsters
-  Shrimp
-  Squid

PLANTS

-  Kelp and Seagrasses
-  Terrestrial Plants

MULTIGROUP

The polygon color and pattern are the same for all the animals in one group. When there is more than one group of animals in one polygon, the polygon is then assigned the multigroup color and pattern. Also associated with each polygon on the map is a number (located under the icon for the polygon). This number references a table on the reverse side of the map with a complete list of species found in the polygon as well as life-history information on each species.

There are some species that are found throughout the nearshore zone on the map. While it is important to note the presence of these species, showing these distributions as polygons would cover large areas. In addition to providing no significant increase in the level of information presented to the user, it would make the maps very difficult to read. In response to this problem, species found in over 25 percent of the water area are identified in a box stating that they are "COMMON THROUGHOUT". This approach informs the user of the presence of these species, while maintaining readability of the map.

BIRDS

Birds are divided into several species subgroups based on behavior and taxonomy. The species table lists all the birds included on the maps sorted by subgroup. These species were included either because of their likelihood of impact by an oil spill or special protection status as threatened or endangered. Bird distribution is shown on the maps as points and polygons. Green dots on the maps depict known nesting sites. Bird polygons are shown as a green hatched pattern; however, if species in addition to birds are in the polygon, a black hatched (multigroup) pattern is used. The number under the icon references a table on the reverse side of the map. In this table, the first column gives the species name, followed by the state (S) and/or federal (F) species designation for endangered (E) or threatened (T) status. The next column provides an estimate of the concentration of species at this site. Concentration is typically indicated as 'HIGH', 'MED', or 'LOW'. These are very subjective values based on local expert opinion on the relative concentrations in the area. If the actual bird counts are available, as for nesting sites, then the actual count will be shown. The species seasonality is shown in the next twelve columns representing the months of the year. If the species is present at that location in a particular month, an 'X' is placed in the month column. The last four columns denote the times for nesting, laying, hatching, and fledging at this site. For many species there is a temporal shift along with the spatial change, so all the temporal information included in the table is specific to the one polygon or point that it references.

FISH

Fish distributions shown on the map represent spawning areas, areas of particularly high concentrations of selected species, and anadromous streams. The species table lists all the fish included on the maps sorted by subgroups. Because these assemblages include many similar species only one icon is used for all fish, instead of one icon for each subgroup as with the other groups. Concentration or spawning areas for fish are shown as polygons on the map. Fish polygons are shown as a blue hatch pattern; however, if species in addition to fish are in the polygon, a black hatched (multigroup) pattern is used. Anadromous fish streams are shown as a thick blue line. Blue icons are associated with both the polygons and the streams. The number under the icon references a table on the reverse side of the map. In this table, the first column gives the species name. The second column denotes whether the species has been designated endangered (E), threatened (T), or special concentration (SC) status on either the state (S) or federal (F) list. The next column provides an estimate of the concentration of species at this site. Concentration is indicated as 'HIGH', 'MED', or 'LOW'. These estimates are very subjective values based on local expert opinion on the relative concentrations in the area. The species seasonality is shown in the next twelve columns, representing the months of the year. If the species is present at that location in a particular month, an 'X' is placed in the month column. The last two columns denote normal times for spawning (all fish) and outmigration (anadromous fish). For many species there is a temporal shift along with the spatial change, so all the temporal information included in the table is specific to the one polygon or line that it references.

MAMMALS

Coastal California has numerous species of marine mammals that potentially may be impacted by an oil spill. Because of the wide diversity in mammals, both behaviorally and physically, the mammals have been divided into subgroups. Each of these subgroups is represented by a different icon. The species table lists all the species of mammals shown on the maps, sorted by subgroup. In addition, there are a few species of terrestrial mammals that might also be impacted. Concentration areas of the more pelagic species (dolphins, porpoises, and whales) are shown on the map, and the general distributions are indicated in the "COMMON THROUGHOUT" box. Mammal distribution on the maps is shown by a brown hatched polygon. However, if species in addition to mammals are included in the polygon, a black hatched (multigroup) polygon is used. The number under the icon references a table on the reverse side of the map. In this table, the first column gives the species name. The second column denotes whether the species has been designated endangered (E) or threatened (T) status on either the state (S) or federal (F) list. The next column provides an estimate of the concentration of species at this site. Concentration is typically indicated as 'HIGH', 'MED', or 'LOW'. These estimates are very subjective values based on local expert opinions about the relative concentrations in the area. In some cases, such as seal or seal lion haulouts, the actual number of animals likely to be present is indicated. The species seasonality is shown in the next twelve columns, representing the months of the year. If the species is present at that location in a particular month, an 'X' is placed in the month column. The last column indicates the most likely dates for birthing by that species. For many species there is a temporal shift along with the spatial change, so all the temporal information included in the table is specific to the one polygon that it references.

SHELLFISH

Shellfish include crustaceans and mollusks and have been divided into several subgroups. The species table lists all the shellfish shown on the maps sorted by subgroup. Species that are commercially or recreationally important, or any species that is threatened or endangered are included. The distribution of shellfish is shown as polygons with an orange hatched pattern. If species in addition to shellfish are included in the polygon, a black hatched (multigroup) pattern is used. Orange icons are associated with the polygons, and the silhouette of the subgroup is shown. The number under the icon references a table on the reverse side of the map. In this table, the first column gives the species name. The second column denotes whether the species has been designated endangered (E) or threatened (T) status on either the state (S) or federal (F) list. The next column provides an estimate of the concentration of species at this site. Concentration is indicated as 'HIGH', 'MED', or 'LOW'. These estimates are very subjective values based on local expert opinions on the relative concentrations in the area. The species seasonality is shown in the next twelve columns, representing the months of the year. If the species is present at that location in a particular month, an 'X' is placed in the month column. The last column indicates dates for spawning. For many species there is a temporal shift along with the spatial change, so all the temporal information included in the table is specific to the one polygon that it references.


PLANTS


The plants are divided into two subgroups: kelp and sea-grasses; and terrestrial plants. The terrestrial plants shown are only those on the state or federal list of threatened or endangered species. Terrestrial plants are seldom directly affected by oil; however, it is possible that cleanup operations might destroy some of the plants or their habitat. The general locations of threatened or endangered plants are shown, so that the appropriate agency can be notified and cleanup and response efforts can be planned accordingly. The species table lists all the plants shown on the


maps. The plants, whether terrestrial or aquatic, are shown as polygons with a purple hatch pattern. If species in addition to plants are present in the polygons, a black hatch (multigroup) pattern is used. Purple icons are associated with the polygons, and the silhouette of the subgroup is shown. The number under the icon references a table on the reverse side of the map. In this table, the first column gives the species name. The second column denotes whether the species has been designated endangered (E) or threatened (T) status on either the state (S) or federal (F) list. The next column provides an estimate of the concentration of species at this site. Concentration is typically indicated as 'HIGH', 'MED', or 'LOW'. These estimates are very subjective values based on local expert opinions on the relative concentrations in the area. The last twelve columns provide information on the plants seasonality. All 12 months are marked with an 'X' since the plants are present all year. This method was used to make tables consistent with those of the other species found on the maps.


HUMAN-USE FEATURES


The human-use features depicted on the maps are those that either could be impacted by an oil spill or could provide access to the cleanup operation. All the features are represented by icons indicating the type of feature. If the icon is not placed on the location of the feature, a leader line is drawn from the icon to the proper location.

 Access—Sites where beach access by vehicle is possible. This information was provided by CDF&G or observed during the overflights.


 Airport—Location of airfields or airports whether they are manned or unmanned. The locations were obtained from visual observations during the overflights or from U.S. Geological Survey (USGS) topographic maps.


 Aquaculture—Location of aquaculture facilities including hatcheries and oyster farms. This information was provided by CDF&G.


 Archaeological site—Location of known archaeological sites in close proximity to the shoreline. This information was provided by CDF&G.


 Boat ramp—Location of boat ramps. This information is from CDF&G, overflight observations, or topographic maps.


 Coast Guard—Location of Coast Guard facilities. This information is from CDF&G and topographical maps.


 Commercial fishing—Areas heavily used for commercial fishing. This information was provided by CDF&G.


 Historical site—Location of all historical sites in close proximity to the shoreline. The information was provided by CDF&G.


 Hoist—Location of facilities which have the capability to hoist boats into and out of the water. This information was provided by CDF&G.


 Marina—Location of any marinas. This information is from CDF&G, overflight observations, or topographic maps.


 Marine sanctuary—The boundaries for the marine sanctuaries were provided by NOAA. The boundaries were entered based on the latitude/longitude point definitions of the marine sanctuary boundaries.


 National park—An icon is used to show the location of the national park, but the digitized boundary was provided by CDF&G.


 State park—An icon is used to show the location of the state park, but the digitized boundary was provided by CDF&G.


 Recreational fishing/boating—General areas where there is heavy recreational fishing or boating. This information was provided by CDF&G.

 Recreational beach—Location of a recreational beach. These sites are indicated with an icon; the beach boundaries were not digitized. Information was provided by CDF&G.

 Reserve, preserve, refuge, or area of special biological significance (ASBS)—All boundaries for the reserves, preserves, refuges, or any other managed and regulated wildlife area were provided by CDF&G. The boundary is shown on the map with an icon and the name in the boundary.

 Subsistence area—Areas utilized by native Americans for collection of subsistence seafoods. Information was provided by CDF&G.

 Village—Native American villages. The boundaries are not delineated. Information was provided by CDF&G.

 Water intakes—Symbol is placed at the location of a water intake. The location information was provided by CDF&G.

For many of these features, the name of the feature, manager/owner, contact, and a phone number were provided. The information is listed below the maps. If at least a name is available for the site, it is included in the list.

Point Cabrillo Lighthouse, State Coastal Conservancy	415/464-1015
Tomales Bay Ecological Reserve, CDF&G	Tom Moore 707/875-2521
Wildlife Refuge, Calif. State Coastal Conservancy	415/464-1015

NAME/OWNER	CONTACT	PHONE
------------	---------	-------

FEDERALLY MANAGED PARKS AND SANCTUARIES

NOAA Sanctuaries and Reserves Division

Cordell Bank National Marine Sanctuary		
Gulf of Farallones National Marine Sanctuary	Ed Ueber	415/556-3509

National Park Service

Point Reyes National Seashore	Bill Shook	415/663-8525
Redwood National Park	Bill Donati	707/464-6101

STATE MANAGED PARKS AND RESERVES

California Department of Parks and Recreation

Chadbourne Gulch State Park		707/937-5804
Del Norte Coast Redwoods State Park	Ken Wilbur	707/445-6547
Greenwood Creek Project		707/937-5804
Humboldt Lagoons State Park		707/488-5435
MacKerricher State Park		707/964-9112
Mendocino Headlands State Park		707/937-5804
Navarro River State Park		707/937-5804
Patrick's Point State Park		707/677-3570
Pelican Beach State Park	Dan Scott	707/464-9533
Prairie Creek Redwoods State Park		707/488-2171
Russian Gulch State Park		707/937-5804
Salt Point State Park	Brian Hickey	707/865-2391
Tomales Bay State Park	Carlos Porrata	415/669-1140
Tomales Bay State Park—Millerton Point	Carlos Porrata	415/669-1140
Van Damme State Park		707/937-5804

Del Norte County Department of Parks and Beaches

Beach Front Park		707/464-7237
Clifford Kamph Memorial Park		707/464-7237

RECREATIONAL BEACHES

California Department of Parks and Recreation

Manchester State Beach		707/937-5804
Schooner Gulch State Beach	David Burtlett	707/937-5804
Sonoma Coast State Beach	Brian Hickey	707/865-2391
Sonoma Coast State Beach—Bodega Head	Brian Hickey	707/865-2391
Sonoma Coast State Beach—Goat Rock	Brian Hickey	707/865-2391
Sonoma Coast State Beach—Russian Gulch	Brian Hickey	707/865-2391
Trinidad State Beach		707/445-4567

Humboldt County Department of Parks

Big Lagoon County Park		707-445-7651
------------------------	--	--------------

Sonoma County Regional Parks

Doran Regional Park	Tom Meyskens	707/875-3540
---------------------	--------------	--------------

Miscellaneous Beaches

Fish Rock Beach/Anchor Bay	Mike Henderson	707/884-4222
Landing State Beach		
Recreational Beach		707/464-9506
Ship Ashore and Salmon Harbor Resorts		707/487-3141
Westport Union		

RESERVES, PRESERVES, AND REFUGES

Bird Rock - ASBS, National Park Service	Bill Shook	415/663-8525
Bodega Marine Life Refuge, University of California	Peter Connors	707/875-2020
Del Mar Landing Ecological Reserve, CDF&G	Mike Williams	707/585-9768
Estero de Limantour Reserve, CDF&G	Tom Moore	707/875-2521
Gerstle Cove ASBS, California Department of Parks and Recreation	Brian Hickey	707/865-2391
Humboldt Bay National Wildlife Refuge	Keven Foerster	707/733-5406
Indian Island, Wiyot Tribal Council		707/773-5055
Jughandle State Reserve, National Park Service		707/937-5804
Kelp Beds at Saunders Reef ASBS, CDF&G	Steve Morse	707/964-0924
King Range National Conservation Area		707/822-7648
Kruse Rhododendron State Reserve		
Mendocino Coast Botanical Garden		707/964-4352
Ocean Cove Reserve	Gary Minaro	707/847-3422
Point Reyes Headland Reserve, CDF&G	Tom Moore	707/875-2521

GEOGRAPHIC INFORMATION SYSTEM DATA

The entire atlas product is stored in digital form in a Geographic Information System (GIS). The information is stored as maps and associated databases. The format for the data varies depending on the type of information or features for which the data are being stored. The three major formats are shoreline habitat classification, biological resources, and human-use features.

Under separate cover are a complete data dictionary, metadata, and descriptive information for the digital data sets and maps that were used to create this atlas. Below is a brief synopsis of the information contained in the digital version. Please refer to the metadata file for full explanations of the data and its structure.

SHORELINE HABITAT CLASSIFICATION

The shoreline habitat classification is stored as lines or polygons with the data identifying the type of habitat associated with the line. In many cases, a shoreline may have two or three different classifications. These multiple classifications are represented on the maps by double and triple lines, and in the database by ESI#1/ESI#2 where ESI#1 is the landward-most classification and ESI#2 is the seaward-most classification.

SENSITIVE BIOLOGICAL RESOURCES

Biological resources are shown on the map by colored and shaded polygons, colored lines and dots, and colored icons. The associated table helps to further identify the resources. In the digital copy, the resources are depicted as lines, points, or polygons. Associated with each map feature is a unique identification number which is linked to a series of databases that further identify the resources. The first data set consists of a list of the species, concentration of each species, and an expert contact for the species. This dataset is then linked to a dataset that describes the life history of each species (temporal presence and reproductive times at month resolution) for the specified map feature. Other databases linked to the first data set are the species identification database, which includes common and scientific names for all species and threatened or endangered status, and the experts database, which includes the name, agency, address, phone number, geographical area of expertise, and biological area of expertise for each of the experts referenced.

HUMAN-USE FEATURES

Human-use features are represented on the maps as an icon describing the feature. In the digital file, the feature location is represented by a point. Attached to the point is a data file that contains the fields for the name of the owner/manager, phone number at which the person can be contacted, identification of the type of feature, and a brief description of the feature. This information is incomplete and may change frequently.

ACKNOWLEDGMENTS

This project was supported jointly by NOAA's Hazardous Materials Response and Assessment Division, Robert Pavia, Project Manager, and the Office of Oil Spill Prevention and Response, Don Lollock, Program Manager. James Morris, Scientific Support Coordinator from NOAA, assisted with many aspects of the logistical arrangements and participated in the field surveys. Don Lollock, Dale Watkins, and Mel Odemar of OSPR's management staff made critical arrangements and participated in some of the field surveys.

All the biological data included on these maps were provided by John Tarpley, Jim Hardwick, Joe Lesh, Melissa Boggs, and Heidi Togstad of CDF&G. They in turn collected the information from numerous people throughout the state of California. The data collection effort was coordinated by Randy Imai of CDF&G.

Kim McCleneghan contributed significantly to the formulation of the project and participated in all phases of the field work. Much of the field work at the ground-truth stations was carried out by personnel from OSPR's Science program, including: Jack Ames, Melissa Boggs, Larry Espinosa, John Grant, Jim Hardwick, Randy Imai, Joe Lesh, Robin Lewis, Dave Rasmussen, John Tarpley, and Heidi Togstad.

At Research Planning, Inc. (RPI), Jeffrey Dahlin was the project biologist and was responsible for the organization of the biological data and the automation of the maps. Shoreline mapping was conducted by Jacqueline Michel and Miles O. Hayes. James Olsen, Scott Johnson, William Holton, Mark White, Lee Diveley, and Niles Shiroff entered the data and produced the final maps. Mike Bise designed the cover, graphics support was provided by Joe Holmes, and Dot Zaino prepared the text.

SPECIES TABLES*

SPECIES TABLES*

Common Name	Species Name
BIRDS	
ALCIDS AND PELAGIC BIRDS	
Ashy storm-petrel	<i>Oceanodroma homochroa</i>
Cassin's auklet	<i>Ptychoramphus aleuticus</i>
Common murre	<i>Uria aalge</i>
Fork-tailed storm-petrel	<i>Oceanodroma furcata</i>
Leach's storm-petrel	<i>Oceanodroma leucorhoa</i>
<u>Marbled murrelet</u>	<i>Brachyramphus marmoratus</i>
Newell's (Manx) shearwater	<i>Puffinus puffinus newelli</i>
Pigeon guillemot	<i>Cephus columba</i>
Pink-footed shearwater	<i>Puffinus creatopus</i>
Sooty shearwater	<i>Puffinus griseus</i>
Tufted puffin	<i>Fratercula cirrhata</i>
DIVING BIRDS	
Brandt's cormorant	<i>Phalacrocorax penicillatus</i>
<u>Brown pelican</u>	<i>Pelecanus occidentalis</i>
Common loon	<i>Gavia immer</i>
Cormorant	<i>Phalacrocorax sp.</i>
Double-crested cormorant	<i>Phalacrocorax auritus</i>
Eared grebe	<i>Podiceps nigricollis</i>
Horned grebe	<i>Podiceps auritus</i>
Pelagic cormorant	<i>Phalacrocorax pelagicus</i>
Pied-billed grebe	<i>Podilymbus podiceps</i>
Red-necked grebe	<i>Podiceps grisegena</i>
Western grebe	<i>Aechmophorus occidentalis</i>
White pelican	<i>Pelecanus erythrorhynchos</i>
GULLS AND TERNS	
Western gull	<i>Larus occidentalis</i>
<u>California least tern</u>	<i>Sterna antillarum browni</i>
Caspian tern	<i>Sterna caspia</i>
RAPTORS	
<u>Bald eagle</u>	<i>Haliaeetus leucocephalus</i>
Northern harrier	<i>Circus cyaneus</i>
Osprey	<i>Pandion haliaetus</i>
<u>Peregrine falcon</u>	<i>Falco peregrinus</i>
SHOREBIRDS	
Black oystercatcher	<i>Haematopus bachmani</i>
Black turnstone	<i>Arenaria melancephala</i>
Black-bellied plover	<i>Pluvialis squatarola</i>
Greater yellowlegs	<i>Tringa melanaleuca</i>
Marbled godwit	<i>Limosa fedoa</i>
Northern phalarope (red-necked)	<i>Phalaropus lobatus</i>
Red phalarope	<i>Phalaropus fulicarius</i>
Western snowy plover	<i>Charadrius alexandrinus nirosus</i>
Willet	<i>Catoptrophorus semipalmatus</i>
TERRESTRIAL, COASTAL, AND/OR MARSH BIRDS	
Bank swallow	<i>Riparia riparia</i>
Black swift	
Saltmarsh common yellowthroat	
WADING BIRDS	
American avocet	<i>Recurvirostra americana</i>
American bittern	<i>Botaurus lentiginosus</i>
Black-crowned night heron	<i>Nycticorax nycticorax</i>
Black-necked stilt	<i>Himantopus mexicanus</i>
<u>California black rail</u>	<i>Laterallus jamaicensis coturniculus</i>
<u>Clapper rail</u>	<i>Rallus longirostris</i>
Great blue heron	<i>Ardea herodias</i>
Great egret	<i>Casmerodius albus</i>
Sora rail	<i>Porzana carolina</i>
Virginia rail	<i>Rallus limicola</i>
WATERFOWL	
<u>Aleutian goose</u>	<i>Branta canadensis leucopareia</i>
American coot	<i>Fulica americana</i>
Black brant	<i>Branta bernicla</i>
Black scoter	<i>Melanitta nigra</i>
Bufflehead	<i>Bucephala albeola</i>
Canada goose	<i>Branta canadensis</i>
Canvasback	<i>Aythya valisineria</i>
Greater scaup	<i>Aythya marila</i>
Green-winged teal	<i>Anas crecca</i>
Harlequin duck	<i>Histrionicus histrionicus</i>
Lesser scaup	<i>Aythya affinis</i>
Northern pintail	<i>Anas acuta</i>
Red-breasted merganser	<i>Mergus serrator</i>
Ruddy duck	<i>Oxyura jamaicensis</i>
Surf scoter	<i>Melanitta perspicillata</i>
Tundra swan	<i>Cygnus columbianus</i>
White-winged scoter	<i>Melanitta deglandi</i>

Common Name	Species Name
MAMMALS	
DOLPHINS	
Bottlenose dolphin	<i>Tursiops truncatus</i>
Dall's porpoise	<i>Phocoenoides dalli dalli</i>
Harbor porpoise	<i>Phocoena phocoena</i>
Northern right-whale dolphin	<i>Lissodelphis borealis</i>
Pacific white-sided dolphin	<i>Lagenorhynchus obliquidens</i>
Risso's dolphin	<i>Grampus griseus</i>
MUSTELIDS AND RODENTS	
River otter	<i>Lutra canadensis</i>
SEA OTTERS	
<u>Sea otter</u>	<i>Enhydra lutris</i>
SEALS AND SEA LIONS	
Harbor seal	<i>Phoca vitulina</i>
Northern elephant seal	<i>Mirounga angustirostris</i>
California sea lion	<i>Zalophus californianus</i>
<u>Northern (Steller) sea lion</u>	<i>Eumetopias jubatus</i>
Northern fur seal	<i>Callorhinus ursinus</i>
WHALES	
<u>Blue whale</u>	<i>Balaenoptera musculus</i>
<u>Fin whale</u>	<i>Balaenoptera physalus</i>
<u>Gray whale</u>	<i>Eschrichtius robustus</i>
<u>Humpback whale</u>	<i>Megaptera novaeangliae</i>
Minke whale	<i>Balaenoptera acutorostrata</i>
<u>Sperm whale</u>	<i>Physeter macrocephalus</i>
FISH	
ANADROMOUS	
Chinook salmon (king) (winter run)	<i>Oncorhynchus tshawytscha</i>
Coho salmon (silver)	<i>Oncorhynchus kisutch</i>
Cutthroat trout	<i>Oncorhynchus clarkii</i>
Rainbow trout (steelhead)	<i>Oncorhynchus mykiss</i>
Salmon fishery (commercial)	
Striped bass	<i>Morone saxatilis</i>
KELP AND SEAGRASS SPAWNERS	
Pacific herring	<i>Clupea harengus pallasii</i>
BEACH SPAWNERS	
California grunion	<i>Leuresthes tenuis</i>
Surf smelt	<i>Hypomesus pretiosus</i>
SPECIAL CONCENTRATIONS	
C-O turbot	<i>Pleuronichthys coenosus</i>
California halibut	<i>Paralichthys californicus</i>
Eulachon	<i>Thaleichthys pacificus</i>
Jacksmelt	<i>Atherinopsis californiensis</i>
Lingcod	<i>Ophiodon elongatus</i>
Night smelt	<i>Spirinchus starksi</i>
Northern anchovy	<i>Engraulis mordax</i>
Pacific lamprey	<i>Entosphenus tridentatus</i>
Prickly sculpin	<i>Cottus asper</i>
Redtail surfperch	<i>Amphistichus rhodoterus</i>
Rockfish	<i>Sebastes spp.</i>
Shiner perch	<i>Cymatogaster aggregata</i>
Starry flounder	<i>Platichthys stellatus</i>
Surfperch	<i>Embiotocidae</i>
Threespine stickleback	<i>Gasterosteus aculeatus</i>
<u>Tidewater goby</u>	<i>Eucyclogobius newberryi</i>
SHELLFISH	
BIVALVES	
Common Pacific littleneck clam	<i>Protothaca staminea</i>
Gaper clam	<i>Tresus nuttallii</i>
Geoduck	<i>Panope generosa</i>
Pismo clam	<i>Tivela stultorum</i>
Razor clam (western)	<i>Siliqua patula</i>
Soft-shell clam	<i>Mya arenaria</i>
Washington clam	<i>Saxidomus nuttallii</i>
Native Pacific oyster	<i>Ostrea lurida</i>
Pacific oyster	<i>Crassostrea gigas</i>
California mussel	<i>Mytilus californianus</i>
Rock scallop	<i>Hinnites multirugosus</i>

* Threatened and endangered species are designated by underlining.

SPECIES TABLES*

Common Name	Species Name
CRABS	
Dungeness crab	<i>Cancer magister</i>
Purple shore crab	<i>Hemigrapsus sp. (nudus?)</i>
Red crab	<i>Cancer productus</i>
Rock crab	<i>Cancer antennarius</i>
Yellow crab	<i>Cancer anthonyi</i>
ECHINODERMS	
Red sea urchin	<i>Strongylocentrotus franciscanus</i>
GASTROPODS AND ABALONE	
California brackish water snail	<i>Tryonia imitator</i>
Abalone	<i>Haliotis sp.</i>
Black abalone	<i>Haliotis cracherodii</i>
Pink abalone	<i>Haliotis corrugata</i>
Red abalone	<i>Haliotis rufescens</i>
LOBSTER	
California rock lobster	<i>Panulirus interruptus</i>
SHRIMP	
Bay ghost shrimp	<i>Callinassa californiensis</i>
SQUID	
Market squid	<i>Loligo opalescens</i>

PLANTS

KELP AND SEAGRASSES	
Bull kelp	<i>Nereocystis luetkeana</i>
Eelgrass	<i>Zostera marina</i>
Giant kelp	<i>Macrocystis pyrifera</i>
Surfgrass	<i>Phyllospadix spp.</i>
TERRESTRIAL PLANTS	
<u>Beach layia</u>	<i>Layia carnosa</i>
Beach spectacle pod	<i>Dithyrea maritima</i>
California suaeda	<i>Suaeda californica</i>
<u>Clover lupine</u>	<i>Lupinus tidestromii</i>
Coastal dunes milkvetch	<i>Astragalus tener titi</i>
<u>Menzies wallflower</u>	<i>Erysimum menziesii</i>
<u>Salt marsh bird's-beak</u>	<i>Cordylanthus maritimus</i>
Sand (Monterey) gilia	<i>Gilia tenuiflora arenaria</i>
Surf thistle	<i>Cirsium rhotophilum</i>

* Threatened and endangered species are designated by underlining.

Shoreline Habitat Descriptions

EXPOSED ROCKY CLIFFS

ESI = 1A and 1C

DESCRIPTION

- The intertidal zone is steep (greater than 30° slope), with very little width.
- Sediment accumulations are uncommon and usually ephemeral (classified as 1A), because waves remove the debris that has slumped from the eroding cliffs.
- Where large boulders have accumulated as talus at the base of the cliff, the shoreline has been classified as 1C.
- This shoreline type is seldom used in combination with another shoreline type, however they are often found interspersed with wave-cut platforms.
- There is strong vertical zonation of intertidal biological communities.
- Species density and diversity vary greatly, but barnacles, snails, mussels, seastars, limpets, sea anemones, shore crabs, polychaetes, and macroalgae are often very abundant.
- They are common throughout northern California, comprising about 7 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Oil is held offshore by wave reflecting off the steep cliffs.
- Any oil that is deposited is rapidly removed from exposed faces.
- The most resistant oil would remain as a patchy band at or above the high-tide line.
- Impacts to intertidal communities are expected to be short-term duration. An exception would be where heavy concentrations of a light refined product came ashore very quickly.

RESPONSE CONSIDERATIONS

- Cleanup is usually not required.
- Access can be difficult and dangerous.



EXPOSED SEAWALLS

ESI = 1B

DESCRIPTION

- Seawalls occur in developed areas to provide protection to residential and industrial developments.
- They are composed of concrete or metal bulkheads.
- Organisms, such as barnacles, mussels, and algae, may be common on the lower levels, whereas biota along the upper intertidal zones is sparse.
- They comprise about 0.5 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Oil would percolate between the joints of the structures.
- Oil would coat the intertidal areas of solid structures.
- Biota would be impacted under heavy accumulations.

RESPONSE CONSIDERATIONS

- High-pressure spraying may be required in order to:
 - remove oil;
 - prepare substrate for recolonization of barnacle and mussel communities;
 - minimize aesthetic damage;
 - prevent the chronic leaching of oil from the structure.

EXPOSED WAVE-CUT PLATFORMS

ESI = 2

DESCRIPTION

- The intertidal zone consists of a flat rock bench of highly variable width.
- The shoreline may be backed by a steep scarp or low bluff.
- There may be a perched beach of sand- to boulder-sized sediments at the base of the scarp.
- The platform surface is irregular and tidal pools are common.
- Small accumulations of gravel can be found in the tidal pools and crevices in the platform.
- These habitats can support large populations of encrusting animals and plants, with rich tidal pool communities. Dominant species include barnacles, snails, mussels, seastars, limpets, sea anemones, shore crabs, and polychaetes.
- They are the most common shoreline type in northern California, representing 23 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Oil will not adhere to the rock platform, but rather be transported across the platform and accumulate along the high-tide line.
- Oil can penetrate in beach sediments, if present.
- Persistence of oiled sediments is usually short-term, except in wave shadows or larger sediment accumulations.

RESPONSE CONSIDERATIONS

- Cleanup is usually not required.
- Where the high-tide area is accessible, it may be feasible to remove heavy oil accumulations and oiled debris.



DESCRIPTION

- These beaches are generally flat, wide, and hard-packed.
- They can occur at the upper intertidal zone on wave-cut platforms.
- Where gravel storm berms occur in the upper intertidal zone, they are also denoted on the maps.
- There can be significant seasonal changes in the beach sediments as well as the width and slope of the beach.
- Upper beach fauna are scarce; lower beach fauna (particularly *Emerita*) can be dense, but are highly variable.
- These beaches are very common, comprising 20 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Light oil accumulations will be deposited as oily swashes or bands along the upper intertidal zone.
- Heavy oil accumulations will cover the entire beach surface; the oil will be lifted off the lower beach with the rising tide.
- Maximum penetration of oil into fine-grained sand is about 10 cm and into medium-grained sand is about 15 cm.
- Burial of oiled layers by clean sand within the first few weeks will be less than 30 cm along the upper beach face.
- Organisms living in the beach may be killed by smothering or lethal oil concentrations in the interstitial water.
- Biological impacts include temporary declines in infaunal populations, which can also affect important shorebird foraging areas.

RESPONSE CONSIDERATIONS

- These beaches are among the easiest beach types to clean.
- Cleanup should concentrate on the removal of oil from the upper swash zone after all oil has come ashore.
- Activity through both oiled and dune areas should be severely limited, to prevent contamination of clean areas.
- Manual cleanup, rather than road graders and front-end loaders, is advised to minimize the volume of sand removed from the shore and requiring disposal.
- All efforts should focus on preventing the mixture of oil deeper into the sediments by vehicular and foot traffic.



COARSE-GRAINED SAND TO GRANULE BEACHES

DESCRIPTION

- These beaches are moderate-to-steep, of variable width, and have soft sediments.
- They commonly occur along beaches at river mouths.
- They are commonly backed by dunes or rocky cliffs along exposed, outer coasts.
- Generally species density and diversity is lower than on fine-grained sand beaches.
- They comprise 6 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- During small spills, oil will be deposited primarily as a band along the high-tide line.
- Under very heavy accumulations, oil may spread across the entire beach face, though the oil will be lifted off the lower part of the beach with the rising tide.
- Penetration of oil into coarse-grained sand can reach 25 cm.
- Burial of oiled layers by clean sand can be rapid, and to depths of 60 cm or more.
- Burial to depths over one meter is possible if the oil comes ashore at the start of a depositional period.
- Biological impacts include temporary declines in infaunal populations, which can also affect important shorebird foraging areas.

RESPONSE CONSIDERATIONS

- Remove oil primarily from the upper swash lines.
- Removal of sediment should be limited to avoid erosion problems.
- Mechanical reworking of the sediment into the surf zone may be used to release the oil without sediment removal.
- Activity in the oiled sand should be limited to prevent mixing oil deeper into the beach.
- Use of heavy equipment for oil/sand removal may result in the removal of excessive amounts of sand; manual cleanup may be more effective.



DESCRIPTION

- Moderately sloping beach composed of a mixture of sand and gravel (less than 80 percent of dominant fraction).
- Because of the mixed sediment sizes, there may be zones of pure sand, pebbles, or cobbles.
- There can be large-scale changes in the sediment distribution patterns depending upon season, because of the transport of the sand fraction offshore during storms.
- Because of sediment desiccation and mobility on exposed beaches, there are low densities of attached animals and plants.
- The presence of attached algae, mussels, and barnacles indicates beaches that are relatively sheltered, with the more stable substrate supporting a richer biota.
- They comprise over 10 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- During small spills, oil will be deposited along and above the high-tide swash.
- Large spills will spread across the entire intertidal area.
- Oil penetration into the beach sediments may be up to 50 cm; however, the sand fraction can be quite mobile, and oil behavior is much like on a sand beach if the sand fraction exceeds about 40 percent.
- Burial of oil may be deep at and above the high-tide line, where oil tends to persist, particularly where beaches are only intermittently exposed to waves.
- In sheltered pockets on the beach, pavements of asphalted sediments can form if there is no removal of heavy oil accumulations, because most of the oil remains on the surface.
- Once formed, these asphalt pavements can persist for many years.
- Oil can be stranded in the coarse sediments on the lower part of the beach, particularly if the oil is weathered or emulsified.



RESPONSE CONSIDERATIONS

- Remove heavy accumulations of pooled oil from the upper beachface.
- All oiled debris should be removed.
- Sediment removal should be limited as much as possible.
- Low-pressure flushing can be used to float oil away from the sediments for recovery by skimmers or sorbents. High-pressure spraying should be avoided because of potential for transporting contaminated finer sediments (sand) to the lower intertidal or subtidal zones.
- Mechanical reworking of oiled sediments from the high-tide zone to the upper intertidal zone can be effective in areas regularly exposed to wave activity (as evidenced by storm berms). However, oiled sediments should not be relocated below the mid-tide zone.
- In-place tilling may be used to reach deeply buried oil layers in the middle zone on exposed beaches.

GRAVEL BEACHES

DESCRIPTION

- Gravel beaches are composed of sediments ranging in size from pebbles to boulders.
- They can be very steep, with multiple wave-built berms forming the upper beach.
- Attached animals and plants are usually restricted to the lowest parts of the beach, where the sediments are less mobile.
- The presence of attached algae, mussels, and barnacles indicates beaches that are relatively sheltered, with the more stable substrate supporting a richer biota.
- They are common adjacent to cliffs and platforms, comprising nearly 7 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Deep penetration and rapid burial of stranded oil is likely on exposed beaches.
- On exposed beaches, oil can be pushed over the high-tide and storm berms, pooling and persisting above the normal zone of wave wash.
- Long-term persistence will be controlled by the depth of penetration versus the depth of routine reworking by storm waves.
- On the more sheltered portions of beaches, formation of asphalt pavements is likely where accumulations are heavy.

RESPONSE CONSIDERATIONS

- Heavy accumulations of pooled oil should be removed quickly from the upper beach.
- All oiled debris should be removed.
- Sediment removal should be limited as much as possible.
- Low- to high-pressure flushing can be used to float oil away from the sediments for recovery by skimmers or sorbents.
- Mechanical reworking of oiled sediments from the high-tide zone to the upper intertidal zone can be effective in areas regularly exposed to wave activity (as evidenced by storm berms). However, oiled sediments should not be relocated below the mid-tide zone.
- In-place tilling may be used to reach deeply buried oil layers in the middle intertidal zone on exposed beaches.



RIPRAP

ESI = 6B

DESCRIPTION

- Riprap structures are composed of cobble- to boulder-sized rock fragments.
- Riprap structures are placed for shoreline protection and inlet stabilization.
- Mid- and low-intertidal zone biota on the riprap may be plentiful and varied.
- Riprap structures are relatively common in northern California, representing 4.5 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Deep penetration of oil between the boulders is likely.
- Oil adheres readily to the rough rock surfaces.
- If oil is left uncleaned, it may cause chronic leaching until the oil asphaltizes.
- Resident fauna and flora may be killed by the oil.

RESPONSE CONSIDERATIONS

- When the oil is fresh and liquid, high-pressure spraying and/or water flooding may be effective, making sure to recover all released oil.
- Heavy and weathered oils are more difficult to remove, requiring scrapping and/or hot-water spraying.
- It may be necessary to remove heavily oiled riprap and replace it.



EXPOSED TIDAL FLATS

ESI = 7

DESCRIPTION

- They are composed primarily of sand and mud.
- The presence of sand indicates that tidal or wind-driven currents and waves are strong enough to mobilize the sediments.
- They are usually associated with another shoreline type on the landward side of the flat.
- They occur in bays and along the lower sections of rivers.
- The sediments usually remain water-saturated, with only the topographically higher ridges drying out during low tide.
- Biological utilization can be very high, with large numbers of infauna, heavy use by birds for roosting and foraging, and use as haulouts for marine mammals.
- In northern California, they comprise nearly 3 percent of the shoreline length.

PREDICTED OIL BEHAVIOR

- Oil does not usually adhere to the surface of exposed tidal flats, but rather moves across the flat and accumulates at the high-tide line.
- Deposition of oil on the flat may occur on a falling tide if concentrations are heavy.
- Oil does not penetrate water-saturated sediments.
- Biological damage may be severe, primarily to infauna, thereby reducing food sources for birds and other predators.

RESPONSE CONSIDERATIONS

- Currents and waves can be very effective in natural removal of the oil.
- Cleanup is very difficult (and possible only during low tides).
- The use of heavy machinery should be restricted to prevent mixing of oil into the sediments.
- On sand flats, oil will be removed naturally from the flat and deposited on the adjacent beaches where cleanup is more feasible.



SHELTERED ROCKY SHORES**ESI = 8A****DESCRIPTION**

- They are bedrock shores of variable slope (from vertical cliffs to wide, rocky ledges) that are sheltered from exposure to most wave and tidal energy.
- The wider shores may have some surface sediments, but the bedrock is the dominant substrate type
- Species density and diversity vary greatly, but barnacles, snails, mussels, seastars, limpets, sea anemones, shore crabs, polychaetes, and macroalgae are often very abundant.
- Sheltered rocky shores are rare in northern California, comprising about 0.3 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- Oil will adhere readily to the rough rocky surface, particularly along the high-tide line, forming a distinct oil band.
- Even on wide ledges, the lower intertidal zone usually stays wet (particularly when algae covered), preventing oil from adhering to the rock surface.
- Heavy and weathered oils can cover the upper zone with little impacts to the rich biological communities of the lower zone.
- Where surface sediments are abundant, oil will penetrate into the crevices formed by the surface rubble and pool at the contact of the sediments and the rock surface.
- Where the rubble is loosely packed, oil will penetrate deeply, causing long-term contamination of the subsurface sediments.

RESPONSE CONSIDERATIONS

- Low- to high-pressure spraying at ambient water temperatures is most effective when the oil is fresh.
- Extreme care must be taken not to spray in the biologically rich lower intertidal zone or when the tidal level reaches that zone.
- Cutting of oiled, attached algae is not recommended; tidal action will eventually float this oil off, so sorbent booms should be deployed.

**SHELTERED MAN-MADE STRUCTURES****ESI = 8B****DESCRIPTION**

- These structures include revetments, seawalls, piers, and docks constructed of impermeable materials such as concrete.
- They are found inside harbors and bays in highly developed areas, sheltered from direct exposure to waves.
- They are uncommon, comprising only 1.2 percent of the shoreline.

PREDICTED OIL BEHAVIOR

- On impermeable surfaces, the oil will form a band at the high-tide line.
- If the oil is not removed, it may cause chronic leaching until the oil hardens into an asphalt deposit.

RESPONSE CONSIDERATIONS

- High-pressure spraying may be required to remove oil for aesthetic reasons and to prevent leaching of oil from the structure.
- Cleanup crews should make sure to recover all released oil.

SHELTERED TIDAL FLATS

ESI=9

DESCRIPTION

- Sheltered tidal flats are composed primarily of silt and clay.
- They are present in calm-water habitats, sheltered from major wave activity, and are frequently fronted by marshes.
- Wave energy is very low, although there may be strong tidal currents on parts of the flat and in channels across the flat.
- The sediments are very soft and cannot support even light foot traffic.
- There can be large populations of clams, worms, and snails.
- Bird life is seasonally abundant.
- Sheltered tidal flats comprise 4 percent of the shoreline length.

PREDICTED OIL BEHAVIOR

- Oil does not usually adhere to the surface of sheltered tidal flats, but rather moves across the flat and accumulates at the high-tide line.
- Deposition of oil on the flat may occur on a falling tide if concentrations are heavy.
- Oil will not penetrate the water-saturated sediments at all.
- In areas of high suspended sediments, sorption of oil can result in deposition of contaminated sediments on the flats.
- Biological damage may be severe.

RESPONSE CONSIDERATIONS

- These are high-priority areas necessitating the use of spill protection devices to limit oil-spill impact; deflection or sorbent booms and open water skimmers should be used.
- Cleanup of the flat surface is very difficult because of the soft substrate and many methods may be restricted.
- Manual operations and deployment of sorbents from shallow-draft boats may be helpful.



MARSHES

ESI = 10

DESCRIPTION

- Marshes are intertidal wetlands containing emergent, herbaceous vegetation.
- Width of the marsh can vary widely, from a narrow fringe to extensive areas.
- They are relatively sheltered from waves and strong tidal currents.
- Resident flora and fauna are abundant with numerous species.
- Bird life is seasonally abundant.
- Marshes are found mostly in major bays, such as Humbolt Bay and Tomales Bay, and at river mouths, such as Ten Mile River and Mad River.
- They comprise about 10 percent of the shoreline length.

PREDICTED OIL BEHAVIOR

- Oil adheres readily to marsh vegetation.
- The band of coating will vary widely, depending upon the tidal stage at the time oil slicks are in the vegetation. There may be multiple bands.
- Large slicks will persist through multiple tidal cycles and coat the entire stem from the high-tide line to the base.
- If the vegetation is thick, heavy oil coating will be restricted to the outer fringe, with penetration and lighter oiling to the limit of tidal influence.
- Medium to heavy oils do not readily adhere or penetrate the fine sediments, but can pool on the surface or in burrows.
- Light oils can penetrate the top few centimeters of sediment and deeply into burrows and cracks (up to one meter).

RESPONSE CONSIDERATIONS

- Under light oiling, the best practice is to let the area recover naturally.
- Heavy accumulations of pooled oil can be removed by vacuum, sorbents, or low-pressure flushing. During flushing, care must be taken to prevent transporting oil to sensitive areas down slope or along shore.
- Cleanup activities should be carefully supervised to avoid vegetation damage.
- Any cleanup activity must not mix the oil deeper into the sediments. Trampling of the roots must be minimized.
- Cutting of oiled vegetation should only be considered when other resources present are at great risk from leaving the oiled vegetation in place.



Northern California Environmental Sensitivity Atlas

(click on the blue text to link to that document)

[Introduction discussion \(INTRO_NC.PDF\)](#)

Explains the purpose and content of the atlas. Lists contacts for parks and reserves in the area, species found in the area, and gives shoreline habitat descriptions.

[Legend of symbols used \(LGND_NC.PDF\)](#)

Explains the symbols used on map pages.

[Index Map \(INDEX_NC.PDF\)](#)

Displays the title page of atlas with map outlining the coverage of each Area map. The coverage **boxes are linked** to the screen resolution Area map pages for quick display online. Just select the box to jump to the associated map page.

[Seasonality Data associated with individual Area Maps \(NCASEASN.PDF\)](#)

References tables for each Area map with a complete list of species found in the numbered area of the map as well as life-history information on each species. **Click page title to link to the associated map**, e.g., ESIMAP 1. **Click map page title to link to the associated seasonality data**, e.g., ENVIRONMENTAL SENSITIVITY INDEX MAP.

Area map pages are in 300-dpi (directory STATE\N_CA\HIRES) and **72-dpi** (directory STATE\N_CA\LOWRES) **resolutions**. Screen (72-dpi) resolution map pages are linked to the coverage boxes on the Index map. **Just select the box to open the associated map page**. The 300-dpi resolution maps are included for high-quality prints of the maps. The map page file names take the form nc(map #) hi.pdf. Map # is the box number shown on the Index map and in the lower right corner of the Area map page.